

lmridge: A Comprehensive R Package for Ridge Regression

by Muhammad Imdad Ullah, Muhammad Aslam, and Saima Altaf

Abstract The ridge regression estimator, one of the commonly used alternatives to the conventional ordinary least squares estimator, avoids the adverse effects in the situations when there exists some considerable degree of multicollinearity among the regressors. There are many software packages available for estimation of ridge regression coefficients. However, most of them display limited methods to estimate the ridge biasing parameters without testing procedures. Our developed package, **lmridge** can be used to estimate ridge coefficients considering a range of different existing biasing parameters, to test these coefficients with more than 25 ridge related statistics, and to present different graphical displays of these statistics.

Introduction

For data collected either from a designed experiment or from an observational study, the ordinary least squares (OLS) method does not provide precise estimates of the effect of any explanatory variable (regressor) when regressors are interdependent (collinear with each other). Consider a multiple linear regression (MLR) model,

$$y = X\beta + \varepsilon, \quad (1)$$

where y is an $n \times 1$ vector of observation on dependent variable, X is known design matrix of order $n \times p$, β is a $p \times 1$ vector of unknown parameters and ε is an $n \times 1$ vector of random errors with mean zero and variance $\sigma^2 I_n$, where I_n is an identity matrix of order n .

The OLS estimator (OLSE) of β is given by

$$\hat{\beta} = (X'X)^{-1}X'y, \quad (2)$$

which depends on characteristics of the matrix $X'X$. If $X'X$ is ill-conditioned (near dependencies among various columns (regressors) of $X'X$ exist) or $\det(X'X) \approx 0$, then the OLS estimates are sensitive to a number of errors, such as non-significant or imprecise regression coefficients (Kmenta, 1980) with wrong sign and non-uniform eigenvalues spectrum. Moreover, the OLS method, can yield high variances of estimates, large standard errors, and wide confidence intervals. Quality and stability of the fitted model may be questionable due to erratic behaviour of the OLSE in case when regressors are collinear.

Researchers may tempt to eliminate regressor(s) causing the problem by consciously removing regressors from the model. However, this method may destroy the usefulness of the model by removing relevant regressor(s) from the model. To control variance and instability of the OLS estimates, one may regularize the coefficients, with some regularization methods such as ridge regression (RR), Liu regression, and Lasso regression methods etc., as alternative to OLS. Computationally, RR suppresses the effects of collinearity and reduces the apparent magnitude of the correlation among regressors in order to obtain more stable estimates of the coefficients than the OLS estimates and it also improves accuracy of prediction (see Hoerl and Kennard, 1970a; Montgomery and Peck, 1982; Myers, 1986; Rawlings et al., 1998; Seber and Lee, 2003; Tripp, 1983, etc.).

There are only a few software programs and R packages capable of estimating and / or testing of ridge coefficients. The design goal of our **lmridge** (Imdad and Aslam, 2018b) is primarily to provide functionality of all possible ridge related computations. The output of our developed package (**lmridge**) is consistent with output of existing software/ R packages. The package, **lmridge** also provides the most complete suite of tools for ordinary RR, comparable to those listed in Table 1. For package development and R documentation, we followed Hadley (2015), Leisch (2008) and R Core Team (2015). The **ridge** package by Moritz and Cule (2017) and **lm.ridge()** from the **MASS** (Venables and Ripley, 2002) also provided guidance in coding.

All available software and R packages mentioned in Table 1 are compared with our **lmridge** package. For multicollinearity detection, NCSS statistical software (NCSS 11 Statistical Software, 2016) computes VIF/TOL, R^2 , eigenvalue, eigenvector, incremental and cumulative percentage of eigenvalues and CN. For RR, ANOVA table, coefficient of variation, plot of residuals vs predicted, histogram and density trace of residuals are also available in NCSS. In SAS (Inc., 2011), collin option in the model statement is used to perform collinearity diagnostics while for remedy of multicollinearity, RR can be performed using a ridge option in proc reg statement. The outVIF option results in

	NCSS	SAS	Stata	StatGraphics	Irnest	Ltsbase	penalized	glmnet	ridge	lmridge
<i>Standardization of regressors</i>										
	✓	✓	✓	✓		✓	✓	✓	✓	✓
<i>Estimation and testing of ridge coefficient</i>										
Estimation	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Testing			✓		✓				✓	✓
SE of coef	✓		✓		✓			✓	✓	✓
<i>Ridge related statistics</i>										
R^2		✓	✓	✓					✓	
adj- R^2			✓		✓				✓	
m-scale & ISRM									✓	
Variance									✓	
Bias ²									✓	
MSE						✓	✓		✓	
F-test			✓						✓	
Shrinkage factor									✓	
CN									✓	
σ^2									✓	
C_k									✓	
DF									✓	
EDF									✓	
Eft									✓	
Hat matrix									✓	
Var-Cov matrix									✓	
VIF	✓			✓				✓	✓	
Residuals	✓		✓	✓		✓	✓		✓	
Ridge fitted					✓	✓			✓	
Predict	✓		✓	✓		✓	✓	✓	✓	
<i>Ridge model selection</i>										
CV & GCV			✓				✓	✓		✓
AIC & BIC										✓
PRESS										✓
<i>Ridge related graphs</i>										
Ridge trace	✓	✓		✓					✓	✓
VIF trace	✓	✓		✓					✓	
Bias, var, MSE					✓				✓	
CV, GCV									✓	
AIC & BIC									✓	
m-scale, ISRM									✓	
DF, RSS, PRESS									✓	

Table 1: Comparison of ridge related software and R packages.

VIF values. For RR, Stata (StataCorp, 2014) has no built-in command, however `ridgereg` add-on is available that performs calculation on scalar k. The `Irnest` package (Dissanayake et al., 2016) computes estimators such as OLS, ordinary RR (ORR), Liu estimator (LE), LE type-1,2,3, Adjusted Liu Estimator (ALTE), and their type-1,2,3 etc. Moreover, `Irnest` provides scalar mean square error (MSE), prediction residual error sum of squares (PRESS) values of some of the estimators. The testing of ridge coefficient is performed only on scalar k, however, for vector of k, function `rid()` of `Irnest` package returns only MSE along with value of biasing parameter used. The function `optimum()` of `Irnest` package can be used to get the optimal scalar MSE and PRESS values (Arumairajan and Wijekoon, 2015). Statgraphics standardizes the dependent variable and computes some statistics for detection of collinearity such as R^2 , adj- R^2 , and VIF. Statgraphics also facilitates to perform RR and computes different RR related statistics such as VIF and ridge trace for different biasing parameter used, R^2 , adj- R^2 and standard error of estimates etc. The `Ltsbase` package (Kan-Kilinc and Alpu, 2013, 2015) computes ridge and Liu estimates based on the least trimmed squares (LTS) method. The MSE value from four regression models can be compared graphically if the argument `plot=TRUE` is passed to the `ltsbase()` function. There are three main functions (i) `ltsbase()` computes the minimum MSE values for six models: OLS, ridge, ridge based on LTS, LTS, Liu, and Liu based on LTS method for sequences of biasing parameters ranging from 0 to 1. If `print=TRUE`, `ltsbase()` prints all the MSEs (along with minimum MSE) for ridge, Liu, and ridge & Liu based on LTS method for the sequence of biasing parameters given by the user, (ii) the `ltsbaseDefault()` function returns the fitted values and residual of the six models (OLS, ridge, Liu, LTS, and ridge & Liu based LTS methods) having minimum MSE, and (iii) the `ltsbaseSummary()` function returns the coefficients and the biasing parameter for the best MSE among the four regression models. The `penalized` package (Goeman et al., 2017) is designed for penalized estimation in generalized linear models. The supported models are linear regression, logistic

regression, Poisson regression and the Cox proportional hazard models. The **penalized** package allows an L1 absolute value ("LASSO") penalty, and L2 quadratic ("ridge") penalty or a combination of the two. It is also possible to have a fused LASSO penalty with L1 absolute value penalty on the coefficients and their differences. The **penalized** package also includes facilities for likelihood, cross-validation and for optimization of the tuning parameter. The **glmnet** package (Friedman et al., 2010) has some efficient procedures for fitting the entire LASSO or elastic-net regularization path for linear regression, logistic and multinomial regression model, Poisson regression and Cox model. The **glmnet** can also be used to fit the RR model by setting alpha argument to zero. The **ridge** package fits linear and also logistic RR models, including functions for fitting linear and logistic RR models for genome-wide SNP data supplied as files names when the data are too big to read into R. The RR biasing parameter is chosen automatically using the method proposed by Cule and De Iorio (2012), however value of biasing parameter can also be specified for estimation and testing of ridge coefficients. The function, **lm.ridge()** from **MASS** only fits linear RR model and returns ridge biasing parameters given by Hoerl and Kennard (1970a) and Venables and Ripley (2002) and vector GCV criterion, given by Golub et al. (1979).

There are other software and R packages that can be used to perform RR analysis such as S-PLUS (S-PLUS, 2008), Shazam (Shazam, 2011) and R packages such as **RXshrink** (Obenchain, 2014), **rrBLUP** (Endelman, 2011), **RidgeFusion** (Price, 2014), **bigRR** (Shen et al., 2013), **lpridge** (Seifert, 2013), **genridge** (Friendly, 2017) and **CoxRidge** (Perperoglou, 2015) etc.

This paper outlines the collinearity detection methods available in the existing literature and uses the **mctest** (Imdad and Aslam, 2018a) package through an illustrative example. To overcome the issues of the collinearity effect on regressors a thorough introduction to ridge regression, properties of the ridge estimator, different methods for selecting values of k , and testing of the ridge coefficients are presented. Finally, estimation of the ridge coefficients, methods of selecting a ridge biasing parameter, testing of the ridge coefficients, and different ridge related statistics are implemented in R within the **lmridge**.

Collinearity detection

Diagnosing collinearity is important to many researchers. It consists of two related but separate elements: (1) detecting the existence of collinear relationship among regressors and (2) assessing the extent to which this relationship has degraded the parameter estimates. There are many diagnostic measures used for detection of collinearity in the existing literature provided by various authors (Belsley et al., 1980; Curto and Pinto, 2011; Farrar and Glauber, 1967; Fox and Weisberg, 2011; Gunst and Mason, 1977; Imdadullah et al., 2016; Klein, 1962; Koutsoyiannis, 1977; Kovács et al., 2005; Marquardt, 1970; Theil, 1971). These diagnostics methods assist in determining whether and where some corrective action is necessary (Belsley et al., 1980). Widely used, and the most suggested diagnostics, are value of pair-wise correlations, variance inflation factor (VIF)/ tolerance (TOL) (Marquardt, 1970), eigenvalues and eigenvectors (Kendall, 1957), CN & CI (Belsley et al., 1980; Chatterjee and Hadi, 2006; Maddala, 1988), Leamer's method (Greene, 2002), Klein's rule (Klein, 1962), the tests proposed by Farrar and Glauber (Farrar and Glauber, 1967), Red indicator (Kovács et al., 2005), corrected VIF (Curto and Pinto, 2011) and Theil's measures (Theil, 1971), (see also Imdadullah et al. (2016)). All of these diagnostic measures are implemented in the R package, **mctest**. Below, we use the Hald dataset (Hald, 1952), for testing collinearity among regressors. We then use the **lmridge** package to compute the ridge coefficients for different ridge related statistics and methods of selection of ridge biasing parameter is also performed. For optimal choice of ridge biasing parameter, graphical representations of the ridge coefficients, vif values, cross validation criteria (CV & GCV), ridge DF, RSS, PRESS, ISRM and m-scale versus used ridge biasing parameter are considered. In addition graphical representation of model selection criteria (AIC & BIC) of ridge regression versus ridge DF is also performed. The Hald data are about heat generated during setting of 13 cement mixtures of 4 basic ingredients and used by Hoerl et al. (1975). Each ingredient percentage appears to be rounded down to a full integer. The data set is already bundled in **mctest** and **lmridge** packages.

Collinearity detection: Illustrative example

```
> library("mctest")
> x <- Hald[, -1]
> y <- Hald[, 1]
> mctest (x, y)
Call:
omcdiag(x = x, y = y, Inter = TRUE, detr = detr, red = red, conf = conf,
theil = theil, cn = cn)
```

Overall Multicollinearity Diagnostics

```
MC Results detection
Determinant |X'X|:      0.0011      1
Farrar Chi-Square:      59.8700      1
Red Indicator:          0.5414      1
Sum of Lambda Inverse:  622.3006      1
Theil's Method:          0.9981      1
Condition Number:       249.5783      1
```

1 --> COLLINEARITY is detected
 0 --> COLLINEARITY is not detected by the test

The results from all overall collinearity diagnostic measures indicate the existence of collinearity among regressor(s). These results do not tell which regressor(s) are reasons of collinearity. The individual collinearity diagnostic measures can be obtained through:

```
> imcdiag(x = x, y = y, all = TRUE)
Call:
imcdiag(x = x, y = y, method = method, corr = FALSE, vif = vif,
tol = tol, conf = conf, cvif = cvif, leamer = leamer, all = all)
```

Individual Multicollinearity Diagnostics

	VIF	TOL	Wi	Fi	Leamer	CVIF	Klein
X1	1	1	1	1	0	0	0
X2	1	1	1	1	1	0	1
X3	1	1	1	1	1	0	0
X4	1	1	1	1	0	0	1

1 --> COLLINEARITY is detected
 0 --> COLLINEARITY is not detected by the test

X1, X2, X3, X4, coefficient(s) are non-significant may be due to multicollinearity

R-square of y on all x: 0.9824

* use method argument to check which regressors may be the reason of collinearity

Results from the most of individual collinearity diagnostics suggest that all of the regressors are the reason for collinearity among regressors. The last line of `imcdiag()` function's output suggests that method argument should be used to check which regressors may be the reason of collinearity among different regressors. For further information about method argument, see the help file of `imcdiag()` function.

Ridge regression analysis

In the seminal work by Hoerl (1959, 1962, 1964) and Hoerl and Kennard (1970b,a) have developed ridge analysis technique that purports the departure of the data from orthogonality. Hoerl (1962) introduced the RR, based on the James-Stein estimator by stating that existence of correlation among regressors can cause errors in estimating regression coefficients when applying the OLS method. The RR is similar to the OLS method however, it shrinks the coefficients towards zero by minimizing the MSE of the estimates, making the RR technique better than the OLSE with respect to MSE, when regressors are collinear with each other. A penalty (degree of bias) is imposed on the size of coefficients in the RR to reduce their variances. However, the expected values of these estimates are not equal to the true values and tend to under estimate the true parameter. Though the ridge estimators are biased but have lower MSE (more precision) than the OLSEs have, less sensitive to sampling fluctuations or model misspecification if number of regressors is more than the number of observations in a data set (i.e., $p > n$), and omitted variables specification bias (Theil, 1957). In summary, the RR procedure is intended to overcome the ill-conditioned situation, and is used to improve the estimation of regression coefficients when regressors are correlated and it also improves the accuracy of prediction (Seber and Lee, 2003). Obtaining the ridge model coefficients ($\hat{\beta}_R$) is relatively straight forward, because the ridge coefficients are obtained by solving a slightly modified form of the OLS method.

The design matrix X in Eq. (1) can be standardized, scaled or centered. Usually, standardization of X matrix is done as described by [Belsley et al. \(1980\)](#) and [Draper and Smith \(1998\)](#), that is, $X_j = \frac{x_{ij} - \bar{x}_j}{\sqrt{\sum(x_{ij} - \bar{x}_j)^2}}$; where $j = 1, 2, \dots, p$ such that $\bar{X}_j = 0$ and $X'_j X_j = 1$, where X_j is the j th column of the matrix X . In this way, the new design matrix (say \tilde{X}) that contains the standardized p columns and the matrix $\tilde{X}' \tilde{X}$ will be correlation matrix of regressors. To avoid complexity of different notations and terms, the centered and scaled design matrix \tilde{X} will be represented by X and centered response variable as y .

The ridge model coefficients are estimated as,

$$\hat{\beta}_{R_k} = (X'X + kI_p)^{-1}X'y, \quad (3)$$

where $\hat{\beta}_{R_k}$ is the vector of standardized RR coefficients of order $p \times 1$ and kI_p is a positive semi-definite matrix added to the $X'X$ matrix. Note that for $k = 0$, $\hat{\beta}_{R_k} = \hat{\beta}_{ols}$.

The addition of constant term k to diagonal element of $X'X$ (in other words addition of kI_p to $X'X$) in Eq. (3) is known as penalty and k is called the biasing or shrinkage parameter. Addition of this biasing parameter guarantees the invertibility of $X'X$ matrix, such that there is always a unique solution $\hat{\beta}_{R_k}$ exists ([Draper and Smith, 1998](#); [Hoerl and Kennard, 1970a](#); [McCallum, 1970](#)) and the condition number (CN) of $X'X + kI$ ($CN_k = \sqrt{\frac{\lambda_1 + k\lambda_p}{\lambda_p + k\lambda_1}}$) also becomes smaller as compared to that of $X'X$, where λ_1 is the largest and λ_p is the smallest eigenvalues of the correlation matrix $X'X$. Therefore, the ridge estimator (RE) is an improvement over the OLSE for collinear data.

It is desirable to select the smallest value of k for which stabilized regression coefficients occur and there always exists a particular value of k for which the total MSE of the REs is less than the MSE of the OLSE, however, the optimum value of k (which produces minimum MSE as compared to other values of k s) varies from one application to another and hence optimal value of k is unknown. Any estimator that has a small amount of bias, less variance and substantially more precise than an unbiased estimator may be preferred since it will have larger probability of being close to the true parameter being estimated. Therefore, criterion of goodness of estimation considered in the RR is the minimum total MSE.

Properties of the ridge estimator

Let X_j denotes the j th column of X ($1, 2, \dots, p$), where $X_j = (x_{1j}, x_{2j}, \dots, x_{nj})'$. As already discussed, assume that the regressors are centered such that $\sum_{i=1}^n x_{ij} = 0$ and $\sum_{i=1}^n x_{ij}^2 = 1$ and the response variable y is centered.

The RR is the most popular among biased methods, because of its relationship to the OLS method and statistical properties of the RE are also well defined. Most of the RR properties have been discussed, proved and extended by many researchers such as [Allen \(1974\)](#); [Hemmerle \(1975\)](#); [Hoerl and Kennard \(1970b,a\)](#); [Marquardt \(1970\)](#); [McDonald and Galarneau \(1975\)](#); [Newhouse and Oman \(1971\)](#). Table 2 lists the RR properties.

Theoretically and practically, the RR is used to propose some new methods for the choice of the biasing parameter k to investigate the properties of RE, since biasing parameter plays a key role while the optimal choice of k is the main issue in this context. In the literature, there are many methods for estimating the biasing parameter k (see [Allen, 1974](#); [Guilkey and Murphy, 1975](#); [Hemmerle, 1975](#); [Hoerl and Kennard, 1970b,a](#); [McDonald and Galarneau, 1975](#); [Obenchain, 1977](#); [Hocking et al., 1976](#); [Lawless and Wang, 1976](#); [Vinod, 1976](#); [Kasarda and Shih, 1977](#); [Hemmerle and Brantle, 1978](#); [Wichern and Churchill, 1978](#); [Nordberg, 1982](#); [Saleh and Kibria, 1993](#); [Singh and Tracy, 1999](#); [Wencheko, 2000](#); [Kibria, 2003](#); [Khafaf and Shukur, 2005](#); [Alkhamisi et al., 2006](#); [Alkhamisi and Shukur, 2007](#); [Khafaf, 2013](#), among many more), however, there is no consensus about which method is preferable ([Chatterjee and Hadi, 2006](#)). Similarly, each of the estimation method of biasing parameter cannot guarantee to give a better k or even cannot give a smaller MSE as compared to that for the OLS.

Methods of selecting values of k

The optimal value of k is one which gives minimum MSE. There is one optimal k for any problem, while a wide range of k ($0 < k < k_{opt}$) give smaller MSE as compared to that of the OLS. For collinear data, a small change in k varies the RR coefficients rapidly. At some values of k , the ridge coefficients get stabilized and the rate of change slow down gradually to almost zero. Therefore, a disciplined way of selecting the shrinkage parameter is required that minimizes the MSE. The biasing parameter k depends on the true regression coefficients (β) and the variance of the residuals σ^2 , unfortunately

sr.#	Property	Formula
1)	Mean	$E(\hat{\beta}_R) = (X'X + kI_p)^{-1}X'\beta$
2)	Shorter regression coeffs.	$\hat{\beta}_R'\hat{\beta}_R \leq \hat{\beta}'\hat{\beta}$
3)	Linear transformation	$\hat{\beta}_R = Z\hat{\beta}$, where $Z = (X'X + kI)^{-1}X'$
4)	Variance	$Var(\hat{\beta}_R) = \sigma^2 \sum_{j=1}^p \frac{\lambda_j}{(\lambda_j+k)^2}$
		$Cov(\hat{\beta}_R) = Cov(Z\hat{\beta})$
5)	Var-Cov matrix	$= \sigma^2(X'X + kI)^{-1}X'X(X'X + kI)^{-1}$ $= \sigma^2[VIF]$ $Bias(\hat{\beta}_R) = -k(X'X + kI)^{-1}\beta$
6)	Bias	$= -k P diag\left(\frac{1}{\lambda_j+k}\right) P'\beta$
7)	MSE	$MSE = \sigma^2 \sum_{j=1}^p \frac{\lambda_j}{(\lambda_j+k)^2} + \sum_{j=1}^p \frac{k^2\alpha_j^2}{(\lambda_j+k)^2}$
8)	Distance between $\hat{\beta}_R$ and β	$\hat{\beta}_R$ and the true vector of β have minimum distance
9)	Inflated RSS	$\phi_0 = k^2\hat{\beta}_R'(X'X)^{-1}\hat{\beta}_R$
10)	R_R^2	$R_R^2 = \frac{\hat{\beta}_R X'y - k\hat{\beta}_R \hat{\beta}_R}{y'y}$
11)	Sampling fluctuations	The $\hat{\beta}_R$ is less sensitive to the sampling fluctuation
12)	Accurate prediction	$\sigma_{f_R}^2 = \sigma^2 \left[1 + x' P diag\left(\frac{\lambda_j}{(\lambda_j+k)^2}\right) P'x \right] + (Bias(\hat{\beta}_R))^2$
13)	Wide range of k	$0 < k < k_{max}$, have smaller set of MSE than OLSE
14)	Optimal k	An optimal k always exists that gives minimum MSE
15)	DF Ridge	$d_{f_R} = EDF = \sum_{j=1}^p \frac{\lambda_j}{\lambda_j+k} = trace[H_{R_k}],$ where $H_{R_k} = X(X'X + kI)^{-1}X'$
16)	Effective no. of parameters	$EP = trace[2H_{R_k} - H_{R_k}H_{R_k}']$
17)	Residual EDF	$REDF = n - trace[2H_{R_k} - H_{R_k}H_{R_k}'] = n - EP$

Table 2: Properties of the ridge estimator.

these are unknown, but they can be estimated from the sample data.

We classified these estimation method as (i) Subjective or (ii) Objective

Subjective methods

In all these methods, the selection of k is subjective or of judgmental nature and provides graphical evidence of the effect of collinearity on the regression coefficient estimates and also accounts for variation by the RE as compared to the OLSE. In these methods, the reasonable choice of k is done using the ridge trace, df trace, VIF trace and plotting of bias, variance, and MSE. The ridge trace is a graphical representation of regression coefficients $\hat{\beta}_R$, as a function of k over the interval $[0, 1]$. The df trace and VIF trace are like the ridge trace plot in which EDF and VIF values are plotted against k . Similarly, plotting of bias, variance, and MSE from the RE may also be helpful in selecting an appropriate value of k . All these graphs can be used for selection of optimal (but judgmental) value of k from horizontal axis to assess the effect of collinearity on each of the coefficients. The effect of collinearity is depressed when value of k increases and all the values of the ridge coefficients, EDF and VIF values decrease and/ or may stabilize after certain value of k . These graphical representations do not provide a unique solution, rather they render a vaguely defined class of acceptable solutions. However, these traces are still useful graphical representations to check for some optimal k .

Objective methods

Suppose, we have set of observations $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ and the RR model as given in Eq. (3). Objective methods, to some extent, are similar to judgmental methods for selection of biasing parameter k , but they require some calculations to obtain these biasing parameters. Table 3 lists widely used methods to estimate the biasing parameter k already available in the existing literature. Table 3

also lists other statistics that can be used for the selection of the biasing parameter k . There are other

method	formula	reference
C_k	$C_k = \frac{SSR_k}{s^2} - n + 2 + 2 \operatorname{trace}(H_{R_k})$ $= \frac{SSR_k}{s^2} + 2(1 + \operatorname{trace}(H_{R_k})) - n$	Kennard (1971); Mallows (1973)
$PRESS_k$	$PRESS_k = \sum_{i=1}^n (y_i - \hat{y}_{(i,-i)_k})^2$ $= \sum_{i=1}^n e_{(i,-i)_k}^2$	Allen (1971, 1974)
CV	$CV_k = n^{-1} \sum_{i=1}^n (y_i - X_j \hat{\beta}_{j_{R_k}})^2$	Delaney and Chatterjee (1986)
GCV	$GCV_k = \frac{SRR_k}{n - (1 + \operatorname{trace}(H_{R_k}))^2}$	Golub et al. (1979)
ISRM	$ISRM_k = \sum_{j=1}^p \left(\frac{\frac{p}{\lambda_j+k}^2}{\sum_{j=1}^p \frac{\lambda_j}{(\lambda_j+k)^2} \lambda_j} - 1 \right)^2$	Vinod (1976)
m-scale	$m = p - \sum_{j=1}^p \frac{\lambda_j}{\lambda_j+k}$	Vinod (1976)
Information criteria	$AIC = n \cdot \log(RSS/n) + 2 \cdot df_{R_k}$ $BIC = n \cdot \log(RSS) + 2 \cdot df_{R_k}$	Akaike (1973); Schwarz (1978)
Effectiveness index (Eft)	$EF = \frac{\sigma^2 \operatorname{trace}(X'X)^{-1} - \sigma^2 \operatorname{trace}(VIF)}{(Bias(\hat{\beta}_R))^2}$	Lee (1979)

Table 3: Objective methods for selection of biasing parameter k .

methods to estimate biasing parameter k . Table 4 lists various methods for the selection of biasing parameter k , proposed by different researchers.

Testing of the ridge coefficients

Investigating of the individual coefficients in a linear but biased regression models such as ridge based, exact and non-exact t type and F test can be used. Exact t -statistics derived by Obenchain (1977) based on the RR for matrix G whose columns are the normalized eigenvectors of $X'X$, is,

$$t^* = \frac{\hat{\beta}_{R_j} - \beta_j}{\sqrt{var(\hat{\beta}_{R_j} - \beta_j)}}, \quad (4)$$

where $j = 1, 2, \dots, p$, $var(\hat{\beta}_{R_j} - \beta_j)$ is an unbiased estimator of the variance of the numerator in Eq. (4), and

$$\beta_j = g_i' \Delta G' [I - (X'X)^{-1} e_i' (e_i (X'X)^{-1} e_i')^{-1}] \hat{\beta}(0),$$

where g_i' is the i th row of G , Δ is the $(p \times p)$ diagonal matrix with i th diagonal element given by $\delta_i = \frac{\lambda_i}{\lambda_i+k}$ and e_i is the i th row of the identity matrix.

It has been established that $\beta_R \sim N(ZX\beta, \phi = Z\Omega Z')$, where $Z = (X'X + kI_p)^{-1}X'$. Therefore, for j th ridge coefficient $\beta_R \sim N(Z_j X\beta, \phi_{jj} = Z_j \Omega Z_j')$ (see Aslam, 2014; Halawa and El-Bassiouni, 2000). Halawa and El-Bassiouni (2000) presented to tackle the problem of testing $H_0 : \beta_j = 0$ by considering a non-exact t type test of the form,

$$t_{R_j} = \frac{\hat{\beta}_{R_j}}{\sqrt{S^2(\hat{\beta}_{R_j})}},$$

where $\hat{\beta}_{R_j}$ is the j th element of RE and $S^2(\hat{\beta}_{R_j})$ is an estimate of the variance of $\hat{\beta}_{R_j}$ given by the i th diagonal element of the matrix $\sigma^2(X'X + kI_p)^{-1}X'X(X'X + kI_p)^{-1}$.

Sr. #	Formula	Reference
1)	$K_{HKB} = \frac{p\hat{\sigma}^2}{\hat{\beta}'\hat{\beta}}$	Hoerl and Kennard (1970a)
2)	$K_{TH} = \frac{(p-2)\hat{\sigma}^2}{\hat{\beta}'\hat{\beta}}$	Thisted (1976)
3)	$K_{LW} = \frac{p\hat{\sigma}^2}{\sum_{j=1}^p \lambda_j \hat{\alpha}_j^2}$	Lawless and Wang (1976)
4)	$K_{DS} = \frac{\hat{\sigma}^2}{\hat{\beta}'\hat{\beta}}$	Dwivedi and Shrivastava (1978)
5)	$K_{LW} = \frac{(p-2)\hat{\sigma}^2 \times n}{\hat{\beta}'X'X\hat{\beta}}$	Venables and Ripley (2002)
6)	$K_{AM} = \frac{1}{p} \sum_{j=1}^p \frac{\hat{\sigma}^2}{\hat{\alpha}_j}$	Kibria (2003)
7)	$\hat{K}_{GM} = \frac{\hat{\sigma}^2}{\left(\prod_{j=1}^p \hat{\alpha}_j^2 \right)^{\frac{1}{p}}}$	Kibria (2003)
8)	$\hat{K}_{MED} = Median\{\hat{\alpha}_j^2\}$	Kibria (2003)
9)	$K_{KM2} = max\left(\frac{1}{\sqrt{\frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}}}\right)$	Muniz and Kibria (2009)
10)	$K_{KM3} = max\left(\sqrt{\frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}}\right)$	Muniz and Kibria (2009)
11)	$K_{KM4} = \left(\prod_{j=1}^p \frac{1}{\sqrt{\frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}}}\right)^{\frac{1}{p}}$	Muniz and Kibria (2009)
12)	$K_{KM5} = \left(\prod_{j=1}^p \sqrt{\frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}}\right)^{\frac{1}{p}}$	Muniz and Kibria (2009)
13)	$K_{KM6} = Median\left(\frac{1}{\sqrt{\frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}}}\right)$	Muniz and Kibria (2009)
14)	$K_{KM8} = max\left(\frac{1}{\sqrt{\frac{\lambda_{max}\hat{\sigma}^2}{(n-p)\hat{\sigma}^2 + \lambda_{max}\hat{\alpha}_j^2}}}\right)$	Muniz et al. (2012)
15)	$K_{KM9} = max\left(\sqrt{\frac{\lambda_{max}\hat{\sigma}^2}{(n-p)\hat{\sigma}^2 + \lambda_{max}\hat{\alpha}_j^2}}\right)$	Muniz et al. (2012)
16)	$K_{KM10} = \left(\prod_{j=1}^p \frac{1}{\sqrt{\frac{\lambda_{max}\hat{\sigma}^2}{(n-p)\hat{\sigma}^2 + \lambda_{max}\hat{\alpha}_j^2}}}\right)^{\frac{1}{p}}$	Muniz et al. (2012)
17)	$K_{KM11} = \left(\prod_{j=1}^p \sqrt{\frac{\lambda_{max}\hat{\sigma}^2}{(n-p)\hat{\sigma}^2 + \lambda_{max}\hat{\alpha}_j^2}}\right)^{\frac{1}{p}}$	Muniz et al. (2012)
18)	$K_{KM12} = Median\left(\frac{1}{\sqrt{\frac{\lambda_{max}\hat{\sigma}^2}{(n-p)\hat{\sigma}^2 + \lambda_{max}\hat{\alpha}_j^2}}}\right)$	Muniz et al. (2012)
19)	$K_{KD} = max\left(0, \frac{p\hat{\sigma}^2}{\hat{\alpha}'\hat{\alpha}} - \frac{1}{n(VIF_j)_{max}}\right)$	Dorugade and Kashid (2010)
20)	$K_{4(AD)} = Harmonic\ Mean[K_i(AD)] \\ = \frac{2p}{\lambda_{max}} \sum_{j=1}^p \frac{\hat{\sigma}^2}{\hat{\alpha}_j^2}$	Dorugade (2014)

Table 4: Different available methods to estimate k .

The statistic t_{R_k} is assumed to follow a Student's t distribution with $(n - p)$ d.f. (Halawa and El-Bassiouni, 2000). Hastie and Tibshirani (1990); Cule and De Iorio (2012) suggested to use $[n - \text{trace}(H_{Rk})]$ d.f. For large sample size, the asymptotic distribution of this statistic is normal (Halawa and El-Bassiouni, 2000). Thus, H_0 is rejected when $|T| > Z_{1-\frac{\alpha}{2}}$.

Similarly, for testing the hypothesis $H_0 : \beta \neq \beta_0$, where β_0 is vector of fixed values. The F statistic for significance testing of the ORR estimator $\hat{\beta}_R$ with $E(\hat{\beta}_R) = ZX\beta$ and estimate of $\text{Cov}(\hat{\beta}_R)$ distributed as $F(DF_{\text{ridge}}, REDF)$ is

$$F = \frac{1}{p} (\hat{\beta}_R - ZX\beta)' (\text{Cov}(\hat{\beta}_R))^{-1} (\hat{\beta}_R - ZX\beta)$$

The R package lmridge

Our R package **lmridge** contains functions related to fitting of the RR model and provides a simple way of obtaining the estimates of RR coefficients, testing of the ridge coefficients, and computation of different ridge related statistics, which prove helpful for selection of optimal biasing parameter k . The package computes different ridge related measures available for the selection of biasing parameter k , and also computes value of different biasing parameters proposed by some researchers in the literature.

The **lmridge** objects contain a set of standard methods such as `print()`, `summary()`, `plot()` and `predict()`. Therefore, inferences can be made easily using `summary()` method for assessing the estimates of regression coefficients, their standard errors, t values and their respective p values. The default function **lmridge** which calls **lmridgeEst()** to perform required computations and estimation for given values of non-stochastic biasing parameter k . The syntax of default function is,

```
lmridge (formula,data,scaling = ("sc","scaled","centered"),K,...)
```

The four arguments of **lmridge()** are described in Table 5:

Argument	Description
formula	Symbolic representation for RR model of the form, response ~ predictors.
data	Contains the variables that have to be used in RR model.
K	The biasing parameter, may be a scalar or vector. If a K value is not provided, $K = 0$ will be used as the default value, i.e., the OLS results will be produced.
scaling	The methods for scaling the predictors. The <code>sc</code> option uses the default scaling of the predictors in correlation form as described in (Belsley, 1991; Draper and Smith, 1998); the <code>scaled</code> option standardizes the predictors having zero mean and unit variance; and the <code>centered</code> option centers the predictors.

Table 5: Description of **lmridge()** function arguments.

The **lmridge()** function returns an object of class "lmridge". The function `summary()`, `kest()`, and `kstats1()` etc., are used to compute and print a summary of the RR results, list of biasing parameter given in Table 4, and ridge related statistics such as estimated squared bias, R^2 and variance etc., after addition of k to diagonal of $X'X$ matrix. An object of class "lmridge" is a list, the components of which are described in Table 6:

Table 7 lists the functions and methods available in **lmridge** package:

The lmridge package implementation in R

The use of **lmridge** is explained through examples by using the Hald dataset.

```
> library("lmridge")
> mod <- lmridge(y ~ X1 + X2 + X3 + X4, data = as.data.frame(Hald),
+ scaling = "sc", K = seq(0, 1, 0.001))
```

The output of linear RR from **lmridge()** function is assigned to an object `mod`. The first argument of the function is `formula`, which is used to specify the required linear RR model for the data provided as second argument. The `print` method for `mod`, an object of class "lmridge", will display the de-scaled coefficients. The output (de-scaled coefficients) from the above command is only for a few selected biasing parameter values.

Object	Description
coef	A named vector of fitted ridge coefficients.
xscale	The scales used to standardize the predictors.
xs	The scaled matrix of predictors.
y	The centered response variable.
Inter	Whether an intercept is included in the model or not.
K	The RR biasing parameter(s).
xm	A vector of means of design matrix X.
rfit	Matrix of ridge fitted values for each biasing parameter k .
d	Singular values of the SVD of the scaled predictors.
div	Eigenvalues of scaled regressors for each biasing parameter k .
scaling	The method of scaling used to standardized the predictors.
call	The matched call.
terms	The terms object used.
Z	A matrix $(X'X + kI_p)^{-1}X'$ for each biasing parameter.

Table 6: Objects from "lmridge" class.

Call:

```
lmridge.default(formula = y ~ ., data = as.data.frame(Hald),
K = seq(0, 1, 0.001))
      Intercept      X1       X2       X3       X4
K=0.01  82.67556 1.31521 0.30612 -0.12902 -0.34294
K=0.05  85.83062 1.19172 0.28850 -0.21796 -0.35423
K=0.5   89.19604 0.78822 0.27096 -0.36391 -0.28064
K=0.9   90.22732 0.65351 0.24208 -0.34769 -0.24152
K=1     90.42083 0.62855 0.23540 -0.34119 -0.23358
```

To get the ridge scaled coefficients mod\$coef can be used,

```
> mod$coef
      K=0.01      K=0.05      K=0.5      K=0.9      K=1
X1  26.800306  24.28399  16.061814  13.316802  12.808065
X2  16.500987  15.55166  14.606166  13.049400  12.689060
X3  -2.862655 -4.83610 -8.074509 -7.714626 -7.570415
X4 -19.884534 -20.53939 -16.272482 -14.004088 -13.543744
```

Objects of class "lmridge" contain components such as rfit, K and coef etc. For fitted ridge model, the generic method summary() is used to investigate the ridge coefficients. The parameter estimates of ridge model are summarized using a matrix of 5 columns namely *estimates*, *estimates (Sc)*, *StdErr (Sc)*, *t values (Sc)* and *P(>|t|)* for ridge coefficients. The following results are shown only for $K = 0.012$ which produces minimum MSE as compared to others values specified in the argument.

```
> summary(mod)
Call:
lmridge.default(formula = y ~ ., data = as.data.frame(Hald), K = 0.012)

Coefficients: for Ridge parameter K= 0.012
            Estimate Estimate (Sc) StdErr (Sc) t-value (Sc) Pr(>|t|)
Intercept    83.1906     -246.5951     269.2195     -0.916  0.3837
X1           1.3046      26.5843      3.8162      6.966  0.0001 ***
X2           0.3017      16.2649      4.6337      3.510  0.0067 ***
X3          -0.1378     -3.0585      3.7655     -0.812  0.4377
X4          -0.3470     -20.1188      4.7023     -4.279  0.0021 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Ridge Summary
      R2    adj-R2 DF ridge      F      AIC      BIC
0.96990  0.95980  3.04587 134.14893 23.24068 58.30578
Ridge minimum MSE= 390.5195 at K= 0.012
P-value for F-test ( 3.04587 , 9.779581 ) = 2.914733e-08
```

Functions	Description
<i>Ridge coefficient estimation and testing</i>	
lmridgeEst()	The main model fitting function for implementation of RR models in R.
coef()	Display de-scaled ridge coefficients.
lmridge()	Generic function and default method that calls lmridgeEst() and returns an object of S3 class "lmridge" with different set of methods to standard generics. It has a print method for display of ridge de-scaled coefficients.
summary()	Standard RR output (coefficient estimates, scaled coefficients estimates, standard errors, t values and p values); returns an object of class "summaryridge" containing the relative summary statistics and has a print method.
<i>Residuals, fitted values and prediction</i>	
predict()	Produces predicted value(s) by evaluating lmridgeEst() in the frame newdata.
fitted()	Displays ridge fitted values for observed data.
residuals()	Displays ridge residuals values.
press()	Generic function that computes prediction residual error sum of squares (PRESS) for ridge coefficients.
<i>Methods to estimate k</i>	
kest()	Displays various k (biasing parameter) values from different authors available in literature and have a print method.
<i>Ridge statistics</i>	
vcov()	Displays associated Var-Cov matrix with matching ridge parameter k values
hatr()	Generic function that displays hat matrix from RR.
infocr()	Generic function that compute information criteria AIC and BIC.
vif()	Generic function that computes VIF values.
rstats1()	Generic function that displays different statistics of RR such as MSE, squared bias and R^2 etc., and have print method.
rstats2()	Generic function that displays different statistics of RR such as df, m-scale and LSRM etc., and have print method.
<i>Ridge plots</i>	
plot()	Ridge and VIF trace plot against biasing parameter k .
bias.plot()	Bias-Variance tradeoff plot. Plot of ridge MSE, bias and variance against k
cv.plot()	Cross validation plots of CV and GCV against biasing parameter k .
info.plot()	Plot of AIC and BIC against k .
isrm.plot()	Plots ISRM and m-scale measure.
rplots.plot()	Miscellaneous ridge related plots such as df-trace, RSS and PRESS plots.

Table 7: Functions and methods in **lmridge** package.

The summary() function also displays ridge related R^2 , adjusted- R^2 , df, F statistics, AIC, BIC and minimum MSE at certain k given in lmridge().

The kest() function, which works with ridge fitted model, computes different biasing parameters developed by researchers, see Table 4. The list of different k values (22 in numbers) may help in deciding the amount of bias needs to be introduced in RR.

```
> kest(mod)
```

Ridge k from different Authors	k values
Thisted (1976):	0.00581
Dwivedi & Srivastava (1978):	0.00291
LW (lm.ridge)	0.05183
LW (1976)	0.00797
HKB (1975)	0.01162
Kibria (2003) (AM)	0.28218

Minimum GCV at	0.01320
Minimum CV at	0.01320
Kibria 2003 (GM):	0.07733
Kibria 2003 (MED):	0.01718
Muniz et al. 2009 (KM2):	14.84574
Muniz et al. 2009 (KM3):	5.32606
Muniz et al. 2009 (KM4):	3.59606
Muniz et al. 2009 (KM5):	0.27808
Muniz et al. 2009 (KM6):	7.80532
Mansson et al. 2012 (KMN8):	14.98071
Mansson et al. 2012 (KMN9):	0.49624
Mansson et al. 2012 (KMN10):	6.63342
Mansson et al. 2012 (KMN11):	0.15075
Mansson et al. 2012 (KMN12):	8.06268
Dorugade et al. 2010:	0.00000
Dorugade et al. 2014:	0.00000

The `rstats1()` and `rstats2()` functions can be used to compute different statistics for a given ridge biasing parameter specified in a call to `lmridge`. The ridge statistics are MSE, squared bias, F statistics, ridge variance, degrees of freedom by [Hastie and Tibshirani \(1990\)](#), condition numbers, PRESS, R^2 , and ISRM etc. Following are the results using `rstats1()` and `rstats2()` functions, for some ($K = 0, 0.012, 0.1, 0.2$).

```
> rstats1(mod)
```

Ridge Regression Statistics 1:

	Variance	Bias^2	MSE	rsigma2	F	R2	adj-R2	CN
K=0	3309.5049	0.0000	3309.5049	5.3182	125.4142	0.9824	0.9765	1376.8806
K=0.012	72.3245	318.1951	390.5195	4.9719	134.1489	0.9699	0.9598	164.9843
K=0.1	19.8579	428.4112	448.2692	5.8409	114.1900	0.8914	0.8552	22.9838
K=0.2	16.5720	476.8887	493.4606	7.6547	87.1322	0.8170	0.7560	12.0804

```
> rstats2(mod)
```

Ridge Regression Statistics 2:

	CK	DF	ridge	EP	REDF	EF	ISRM	m	scale	PRESS
K= 0	6.0000	4.0000	4.0000	9.0000	0.0000	3.9872	0.0000	110.3470		
K= 0.012	4.8713	3.0459	3.2204	9.7796	10.1578	3.6181	0.9541	92.8977		
K= 0.1	4.2246	2.5646	2.9046	10.0954	7.6829	2.8471	1.4354	121.2892		
K= 0.2	3.8630	2.2960	2.7290	10.2710	6.9156	2.5742	1.7040	162.2832		

The residuals, fitted values from the RR and predicted values of the response variable y can be computed using functions `residual()`, `fitted()` and `predict()`, respectively. To obtain the *Var-Cov* matrix, VIF and Hat matrix, the function `vcov()`, `vif()` and `hatr()` can be used. The df are computed by following [Hastie and Tibshirani \(1990\)](#). The results for VIF, *Var-Cov* and diagonal elements of the hat matrix from `vif()`, `vcov()` and `hatr()` functions are given below for $K = 0.012$.

```
> hatr(mod)
> hatr(mod)[[2]]
> diag(hatr(mod)[[2]])
> diag(hatr(lmridge(y ~ ., as.data.frame(Hald), K = c(0, 0.012)))[[2]])
1      2      3      4      5      6      7      8      9      10     11 
0.39680 0.21288 0.10286 0.16679 0.24914 0.04015 0.28424 0.30163 0.12502 0.58426 0.29625 
12      13 
0.12291 0.16294
```

```
> vif(mod)
          X1        X2        X3        X4
k=0    38.49621 254.42317 46.86839 282.51286
k=0.012 2.92917 4.31848 2.85177 4.44723
k=0.1   1.28390 0.51576 1.20410 0.39603
k=0.2   0.78682 0.34530 0.75196 0.28085
```

```
R> vcov(mod)
$`K=0.012`
```

	X1	X2	X3	X4
X1	14.563539	1.668783	11.577483	4.130232
X2	1.668783	21.471027	3.066958	19.075274
X3	11.577483	3.066958	14.178720	4.598000
X4	4.130232	19.075274	4.598000	22.111196

Following are possible uses of some functions to compute different ridge related statistics. For detail description of these functions/ commands, see the **lmridge** package documentation.

```
> mod$rfit
> resid(mod)
> fitted(mod)
> infocr(mod)
> press(mod)
```

For given values of X , such as for first five rows of X matrix, the predicted values for some $K = 0, 0.012, 0.1$, and 0.2 will be computed by predict():

```
> predict(mod, newdata = as.data.frame(Hald[1 : 5, -1]))
      K=0   K=0.012   K=0.1   K=0.2
1 78.49535 78.52225 79.75110 80.73843
2 72.78893 73.13500 74.32678 75.38191
3 105.97107 106.39639 106.04958 105.62451
4 89.32720 89.48443 89.52343 89.65432
5 95.64939 95.73595 96.56710 96.99781
```

The model selection criteria's of AIC and BIC can be computed using infocr() function for each value of K used in argument of ridge(). For some $K = 0, 0.012, 0.1$, and 0.2 , the AIC and BIC values are:

```
> infocr(mod)
      AIC      BIC
K=0    24.94429 60.54843
K=0.012 23.24068 58.30578
K=0.1    24.78545 59.57865
K=0.2    27.98813 62.62961
```

The effect of multicollinearity on the coefficient estimates can be identified by using different graphical displays such as ridge, VIF and df traces, plotting of RSS against df, PRESS vs k , and the plotting of bias, variance, and MSE against K etc. Therefore, for selection of optimal k using subjective (judgmental) methods, different plot functions are also available in **lmridge** package. For example, the ridge (Figure 1) or vif trace (Figure 2) can be plotted using plot() function. The argument to plot functions are abline = TRUE, and type = c("ridge", "vif"). By default, ridge trace will be plotted having horizontal line parallel to horizontal axis at $y = 0$ and vertical line on x -axis at k having minimum GCV.

```
> mod <- lmridge(y ~ ., data = as.data.frame(Hald), K = seq(0, 0.5, 0.001))
> plot(mod)
> plot(mod, type = "vif", abline = FALSE)
> plot(mod, type = "ridge", abline = TRUE)

> bias.plot(mod, abline = TRUE)
> info.plot(mod, abline = TRUE)

> cv.plot(mod, abline = TRUE)
```

The vertical lines in ridge trace and VIF trace suggest the optimal value of biasing parameter k selected at which GCV is minimum. The horizontal line in ridge trace is reference line at $y = 0$ for ridge coefficient against vertical axis.

The bias-variance tradeoff plot (Figure 3) may be used to select optimal k using bias.plot() function. The vertical line in bias-variance tradeoff plot shows the value of biasing parameter k and horizontal line shows minimum MSE for ridge.

The plot of model selection criteria AIC and BIC for choosing optimal k (Figure 4), info.plot() function may be used,

Function cv.plot() plots the CV and GCV cross validation against biasing parameter k for the optimal selection of k (see Figure 5), that is,




Figure 1: Ridge trace plot.




Figure 2: VIF trace.

```
> isrm.plot(mod)
```

The m -scale and ISRM (Figure 6) measures by Vinod (1976) can also be plotted from function of `isrm.plot()` and can be used to judge the optimal value of k .

Function `rplots.plot()` plots the panel of three plots namely (i) df trace, (ii) RSS vs k and (iii) PRESS vs k and may be used to judge the optimal value of k , see Figure 7.

```
> rplots.plot(mod)
```




Figure 3: Bias-variance trade-off.




Figure 4: Information criteria plot (AIC and BIC).

Summary

Our developed **lmridge** package provides the most complete suite of tools for RR available in R, comparable to those available as listed in Table 1. We have implemented functions to compute the ridge coefficients, testing of these coefficients, computation of different ridge related statistics and computation of the biasing parameter for different existing methods by various authors (see Table 4).

We have greatly increased the ridge related statistics and different graphical methods for the selection of biasing parameter k through **lmridge** package in R.

Up to now, a complete suite of tools for RR was not available for an open source or paid version of statistical software packages, resulting in reduced awareness and use of developed ridge related statistics. The package **lmridge** provides a complete open source suite of tools for the computation of




Figure 5: Cross-validation plots (CV and GCV).




Figure 6: m-scale and ISRM plot.

ridge coefficients estimation, testing and computation of different statistics. We believe the availability of these tools will lead to increase utilization and better ridge related practices.

Bibliography

- H. Akaike. A New Look at the Statistical Model Identification. *IEEE Transaction on Automatic Control*, 9 (6):716–723, 1973. URL <https://doi.org/10.1109/TAC.1974.1100705>. [p332]
- M. A. Alkhamisi and G. Shukur. A Monte Carlo Study of Recent Ridge Parameters. *Communications in Statistics - Theory and Methods*, 33(10):2533–2548, 2004. URL <https://doi.org/10.1081/STA-120037300>. [p332]




Figure 7: Miscellaneous ridge plots.

tions in Statistics-Simulation and Computation, 36(3):535–547, 2007. URL <https://doi.org/10.1080/03610910701208619>. [p330]

M. A. Alkhamisi, G. Khalaf, and G. Shukur. Some Modifications for Choosing Ridge Parameter. *Communications in Statistics-Theory and Methods*, 35(11):2005–2020, 2006. URL <https://doi.org/10.1080/03610920600762905>. [p330]

D. M. Allen. Mean Square Error of Prediction as a Criterion for Selecting Variables. *Technometrics*, 13(3):469–475, 1971. URL <http://doi.org/10.2307/1267161>. [p332]

D. M. Allen. The Relationship Between Variable Selection and Data Augmentation and Method for Prediction. *Technometrics*, 16(1):125–127, 1974. URL <http://doi.org/10.2307/1267500>. [p330, 332]

S. Arumairajan and P. Wijekoon. Optimal generalized biased estimator in linear regression model. *Open Journal of Statistics*, 5:403–411, 2015. URL <https://doi.org/10.4236/ojs.2015.55042>. [p327]

M. Aslam. Using Heteroscedasticity-Consistent Standard Errors for the Linear Regression Model with Correlated Regressors. *Communications in Statistics-Simulation and Computation*, 43(10):2353–2373, 2014. URL <https://doi.org/10.1080/03610918.2012.750354>. [p332]

D. A. Belsley. A Guide to Using the Collinearity Diagnostics. *Computer Science in Economics and Management*, 4(1):33–50, 1991. URL <https://doi.org/10.1007/BF00426854>. [p334]

D. A. Belsley, E. Kuh, and R. E. Welsch. *Diagnostics: Identifying Influential Data and Sources of Collinearity*. John Wiley & Sons, New York, 1980. chap. 3. [p328, 330]

S. Chatterjee and A. S. Hadi. *Regression Analysis by Example*. John Wiley & Sons, 4th edition, 2006. [p328, 330]

E. Cule and M. De Iorio. A Semi-Automatic Method to Guide the Choice of Ridge Regression. *Annals of Applied Statistics*, arxiv:1205.0686v1, 2012. URL <https://arxiv.org/abs/1205.0686v1>. [p328, 334]

J. D. Curto and J. C. Pinto. The Corrected VIF (CVIF). *Journal of Applied Statistics*, 38(7):1499–1507, 2011. URL <https://doi.org/10.1080/02664763.2010.505956>. [p328]

N. J. Delaney and S. Chatterjee. Use of the Bootstrap and Cross-Validation in Ridge Regression. *Journal of Business & Economic Statistics*, 4(2):255–262, 1986. URL <http://doi.org/10.2307/1391324>. [p332]

- A. Dissanayake, P. Wijekoon, and R-Core. lmrest: Different Types of Estimators to Deal with Multicollinearity, 2016. URL <https://cran.r-project.org/package=lmrest.R>. package version 3.0. [p327]
- A. V. Dorugade. New Ridge Parameters for Ridge Regression. *Journal of the Association of Arab Universities for Basic and Applied Sciences*, 15:94–99, 2014. URL <https://doi.org/10.1016/j.jaubas.2013.03.005>. [p333]
- A. V. Dorugade and D. N. Kashid. Alternative Method for Choosing Ridge Parameter for Regression. *Applied Mathematical Sciences*, 4(9):447–456, 2010. [p333]
- N. R. Draper and H. Smith. *Applied Regression Analysis*. John Wiley & Sons, New York, 2nd edition, 1998. [p330, 334]
- T. D. Dwivedi and V. K. Shrivastava. On the Minimum Mean Square Error Estimators in a Regression Model. *Communications in Statistics-Theory and Methods*, 7(5):487–494, 1978. URL <https://doi.org/10.1080/03610927808827642>. [p333]
- J. B. Endelman. Ridge Regression and Other Kernels for Genomic Selection with R Package rrBLUP. *Plant Genome*, 4(3):250–255, 2011. URL <http://doi.org/10.3835/plantgenome2011.08.0024>. [p328]
- D. E. Farrar and R. R. Glauber. Multicollinearity in Regression Analysis: The Problem Revisited. *The Review of Economics and Statistics*, 49(1):92–107, 1967. URL <http://doi.org/10.2307/1937887>. [p328]
- J. Fox and S. Weisberg. *An R Companion to Applied Regression*. Sage, Thousand Oaks, CA, 2nd edition, 2011. URL <https://socialsciences.mcmaster.ca/jfox/Books/Companion>. [p328]
- J. Friedman, T. Hastie, and R. Tibshirani. Regularization Paths for Generalized Linear Models via Coordinate Descent. *Journal of Statistical Software*, 33(1):1–22, 2010. URL <http://doi.org/10.18637/jss.v033.i01>. [p328]
- M. Friendly. genridge: Generalized Ridge Trace Plots for Ridge Regression, 2017. URL <https://cran.r-project.org/web/packages/genridge>. R package version 0.6-6. [p328]
- J. J. Goeman, R. J. Meijer, and N. Chaturvedi. Penalized: L1 (Lasso and Fused Lasso) and L2 (Ridge) Penalized Estimation in GLMs and in the Cox Model, 2017. URL <https://cran.r-project.org/web/packages/penalized>. R package version 0.9-50. [p327]
- G. H. Golub, M. Heath, and G. Wahba. Generalized Cross Validation as a Method for Choosing a Good Ridge Parameter. *Technometrics*, 21(2):215–223, 1979. URL <http://doi.org/10.2307/1268518>. [p328, 332]
- W. H. Greene. *Econometric Analysis*. Prentic Hall, New Jersey, 5th edition, 2002. [p328]
- D. K. Guilkey and J. L. Murphy. Direct Ridge Regression Techniques in Cases of Multicollinearity. *Journal of American Statistical Association*, 70(352):769–775, 1975. URL <http://doi.org/10.1080/01621459.1975.10480301>. [p330]
- R. F. Gunst and R. L. Mason. Advantages of Examining Multicollinearities in Regression Analysis. *Biometrics*, 33(1):249–260, 1977. [p328]
- W. Hadley. *R Packages: Organize, Test, Document, and Share Your Code*. O'Reilly Media, 2015. [p326]
- A. M. Halawa and M. Y. El-Bassiouni. Tests of Regression Coefficients Under Ridge Regression Models. *Journal of Statistical-Computation and Simulation*, 65(1–4):341–356, 2000. URL <https://doi.org/10.1080/00949650008812006>. [p332, 334]
- A. Hald. *Statistical Theory with Engineering Applications*. John Wiley & Sons, 1952. [p328]
- T. Hastie and R. Tibshirani. *Generalized Additive Models*. Chapman & Hall, 1990. [p334, 337]
- W. J. Hemmerle. An Explicit Solution for Generalized Ridge Regression. *Technometrics*, 17(3):309–314, 1975. URL <http://doi.org/10.2307/1268066>. [p330]
- W. J. Hemmerle and T. F. Brantle. Explicit and Constrained Generalized Ridge Estimation. *Technometrics*, 20(2):109–120, 1978. URL <http://doi.org/10.2307/1268701>. [p330]
- R. R. Hocking, F. M. Speed, and M. J. Lynn. A Class of Biased Estimators in Linear Regression. *Technometrics*, 18(4):425–437, 1976. URL <http://doi.org/10.2307/1268658>. [p330]

- A. E. Hoerl. Optimum Solution of Many Variables Equations. *Chemical Engineering Progress*, 55:67–78, 1959. [p329]
- A. E. Hoerl. Application of Ridge Analysis to Regression Problems. *Chemical Engineering Progress*, 58: 54–59, 1962. URL <http://doi.org/10.1002/sim.4780030311>. [p329]
- A. E. Hoerl. Ridge Analysis. *Chemical Engineering Progress Symposium Series*, 60:67–77, 1964. [p329]
- A. E. Hoerl and R. W. Kennard. Ridge Regression: Biased Estimation of Nonorthogonal Problems. *Technometrics*, 12(1):55–67, 1970a. URL <http://doi.org/10.2307/1267351>. [p326, 328, 329, 330, 333]
- A. E. Hoerl and R. W. Kennard. Ridge Regression: Application to Nonorthogonal Problems. *Technometrics*, 12(1):69–82, 1970b. URL <http://doi.org/10.2307/1267352>. [p329, 330]
- A. E. Hoerl, R. W. Kennard, and K. F. Baldwin. Ridge Regression: Some Simulations. *Communications in Statistics*, 4(2):105–123, 1975. URL <https://doi.org/10.1080/03610927508827232>. [p328]
- M. U. Imdad and M. Aslam. mctest: An R Package for Detection of Collinearity Among Regressors, 2018a. URL <https://cran.r-project.org/web/packages/mctest/>. R package version 1.2. [p328]
- M. U. Imdad and M. Aslam. lmridge: Linear Ridge Regression with Ridge Penalty and Ridge Statistics, 2018b. URL <https://cran.r-project.org/package=lmridge>. R package version 1.2. [p326]
- M. Imdadullah, M. Aslam, and S. Altaf. mctest: An R Package for Detection of Collinearity Among Regressors. *The R Journal*, 8(2):495–505, 2016. URL <https://journal.r-project.org/archive/accepted/imdadullah-aslam-altaf.pdf>. [p328]
- S. I. Inc. The SAS System for Windows: Release 9.2, 2011. SAS Inst., Cary, NC. [p326]
- B. Kan-Kilinc and O. Alpu. ltsbase: Ridge and Liu Estimates Based on LTS (Least Trimmed Squares) Method, 2013. URL <https://CRAN.R-project.org/package=ltsbase>. R package version 1.0.1. [p327]
- B. Kan-Kilinc and O. Alpu. Combining Some Biased Estimation Methods with Least Trimmed Squares Regression and Its Application. *Revista Colombiana de Estadística*, 38(2):485–502, 2015. URL <http://dx.doi.org/10.15446/rce.v38n2.51675>. [p327]
- J. D. Kasarda and W. F. Shih. Optimal Bias in Ridge Regression Approaches to Multicollinearity. *Sociological Methods and Research*, 5(4):461–470, 1977. URL <https://doi.org/10.1177/004912417700500405>. [p330]
- M. G. Kendall. *A Course in Multivariate Analysis*. Griffin, London, 1957. pp. 70–75. [p328]
- R. W. Kennard. A Note on the Cp-Statistics. *Technometrics*, 13(4):899–900, 1971. URL <https://doi.org/10.1080/00401706.1971.10488863>. [p332]
- G. Khalaf. A Comparison Between Biased and Unbiased Estimators. *Journal of Modern Applied Statistical Methods*, 12(2):293–303, 2013. URL <http://10.22237/jmasm/1383279360>. [p330]
- G. Khalaf and G. Shukur. Choosing Ridge Parameter for Regression Problems. *Communications in Statistics-Theory and Methods*, 34(5):1177–1182, 2005. URL <https://doi.org/10.1081/STA-200056836>. [p330]
- B. M. G. Kibria. Performance of Some New Ridge Regression Estimators. *Communications in Statistics-Simulation and Computation*, 32(2):419–435, 2003. URL <https://doi.org/10.1081/SAC-120017499>. [p330, 333]
- L. R. Klein. *An Introduction to Econometrics*. Prentic Hall, Englewood Cliffs, N. J., 1962. pp. 101. [p328]
- J. Kmenta. *Elements of Econometrics*. Macmillan Publishing Company, New York, 2nd edition, 1980. pp. 431. [p326]
- A. Koutsoyiannis. *Theory of Econometrics*. Macmillan Education Limited, 1977. [p328]
- P. Kovács, T. Petres, and Tóth. A New Measure of Multicollinearity in Linear Regression Models. *International Statistical Review / Revue Internationale de Statistique*, 73(3):405–412, 2005. URL <http://doi.org/10.1111/j.1751-5823.2005.tb00156.x>. [p328]
- J. F. Lawless and P. Wang. A Simulation Study of Ridge and Other Regression Estimators. *Communications in Statistics-Theory and Methods*, 5(4):307–323, 1976. URL <https://doi.org/10.1080/03610927608827353>. [p330, 333]

- W. F. Lee. Model Estimation Using Ridge Regression with the Variance Normalization Criterion. Master thesis, Department of Educational Foundation, Memorial University of Newfoundland, 1979. [p322]
- F. Leisch. *Creating R Packages: A Tutorial*. Compstat 2008-Proceedings in Computational Statistics, Physica Verlage, Heidelberg, Germany, 2008, 2008. URL <ftp://cran.r-project.org/pub/R/doc/contrib/Leisch-CreatingPackages.pdf>. [p326]
- G. S. Maddala. *Introduction to Econometrics*. Macmillan Publishing Company, New York, 1988. [p328]
- C. L. Mallows. Some Comments on Cp. *Technometrics*, 15(4):661–675, 1973. URL <http://doi.org/10.2307/1267380>. [p322]
- D. W. Marquardt. Generalized Inverses, Ridge Regression, Biased Linear Estimation, and Nonlinear Estimation. *Technometrics*, 12(3):591–612, 1970. URL <http://doi.org/10.2307/1267205>. [p328, 330]
- B. T. McCallum. Artificial Orthogonalization in Regression Analysis. *Review of Economics and Statistics*, 52(1):110–113, 1970. URL <http://doi.org/10.2307/1927606>. [p330]
- G. C. McDonald and D. I. Galarneau. A Monte Carlo Evaluation of Some Ridge-Type Estimators. *Journal of the American Statistical Association*, 70(350):407–416, 1975. URL <http://doi.org/10.2307/2285832>. [p330]
- D. C. Montgomery and E. A. Peck. *Introduction to Linear Regression Analysis*. John Wiley & Sons, New York, 1982. [p326]
- S. Moritz and E. Cule. ridge: Ridge Regression with Automatic Selection of the Penalty Parameter, 2017. URL <https://cran.r-project.org/web/packages/ridge>. R package version 2.2. [p326]
- G. Muniz and B. M. G. Kibria. On Some Ridge Regression Estimators: An Empirical Comparisons. *Communications in Statistics-Simulation and Computation*, 38(3):621–630, 2009. URL <https://doi.org/10.1080/03610910802592838>. [p333]
- G. Muniz, B. M. G. Kibria, K. Mansson, and G. Shukur. On Developing Ridge Regression Parameters: A Graphical Investigation. *Statistics and Operations Research Transactions*, 36(2):115–138, 2012. [p333]
- R. H. Myers. *Classical and Modern Regression with Application*. PWS-KENT Publishing Company, 2 edition, 1986. [p326]
- NCSS 11 Statistical Software. Ridge Regression, 2016. URL http://ncss.wpengine.netdna-cdn.com/wp-content/themes/ncss/pdf/Procedures/NCSS/Ridge_Regression.pdf. NCSS, LLC. Kaysville, Utah, USA. [p326]
- J. P. Newhouse and S. D. Oman. An Evaluation of Ridge Estimators. Rand Report, R-716-PR, 1971. [p330]
- L. Nordberg. A Procedure for Determination of a Good Ridge Parameter in Linear Regression. *Communications in Statistics-Simulation and Computation*, 11(3):285–309, 1982. URL <https://doi.org/10.1080/03610918208812264>. [p330]
- B. Obenchain. RXshrink: Maximum Likelihood Shrinkage via Generalized Ridge or Least Angle Regression, 2014. URL <https://cran.r-project.org/web/packages/RXshrink>. R package version 1.0-8. [p328]
- R. L. Obenchain. Classical F-Tests and Confidence Regions for Ridge Regression. *Technometrics*, 19(4):429–439, 1977. URL <http://doi.org/10.2307/1267882>. [p330, 332]
- A. Perperoglou. CoxRidge: Cox Models with Dynamic Ridge Penalties, 2015. URL <https://cran.r-project.org/web/packages/CoxRidge>. R package version 0.9.2. [p328]
- B. S. Price. RidgeFusion: R Package for Ridge Fusion in Statistical Learning, 2014. URL <https://cran.r-project.org/web/packages/RidgeFusion>. R package version 1.0-3. [p328]
- R Core Team. *Writing R Extensions*. R Foundation for Statistical Computing, 2015. Version R 3.2.3. [p326]
- J. O. Rawlings, S. G. Pantula, and D. A. Dickey. *Applied Regression Analysis: A Research Tool*. Springer-Verlag, New York, 2nd edition, 1998. [p326]
- S-PLUS. Insightful Corporation, Seattle, Wt., 2008. URL <http://www.insightful.com>. [p328]

- A. K. M. E. Saleh and B. M. G. Kibria. Performance of Some New Preliminary Test Ridge Regression Estimators and Their Properties. *Communications in Statistics-Theory and Methods*, 22(10):2747–2764, 1993. URL <https://doi.org/10.1080/03610929308831183>. [p330]
- G. E. Schwarz. Estimating the Dimension of a Model. *The Annals of Statistics*, 6(2):461–464, 1978. URL <https://doi.org/10.1214/aos/1176344136>. [p332]
- G. A. F. Seber and A. J. Lee. *Linear Regression Analysis*. John Wiley & Sons, New Jersey, 2 edition, 2003. [p326, 329]
- B. Seifert. lpridge: Local Polynomial (Ridge) Regression, 2013. URL <https://cran.r-project.org/web/packages/lpridge>. R package version 1.0-7. [p328]
- Shazam. *Reference Manual Version 11*, 2011. ISBN: 978-0-9570475-0-1. [p328]
- X. Shen, M. Alam, F. Fikse, and L. Rönnegård. A Novel Generalized Ridge Regression Method for Quantitative Genetics. *Genetics*, 193(4):1255–1268, 2013. URL <http://doi.org/10.1534/genetics.112.146720>. [p328]
- S. Singh and D. S. Tracy. Ridge-Regression Using Scrambled Responses. *Metrika*, LVII(1-2):147–157, 1999. [p330]
- StataCorp. Stata Statistical Software: Release 8, 2014. URL <http://www.stata.com>. College Station, TX: StataCorp LP. [p327]
- H. Theil. Specification Errors and the Estimation of Economic Relationships. *International Statistical Institute (ISI)*, 25(1/3):41–51, 1957. URL <http://doi.org/10.2307/1401673>. [p329]
- H. Theil. *Principles of Econometrics*. John Wiley & Sons, New York, 1971. [p328]
- R. A. Thisted. Ridge Regression, Minimax Estimation, and Empirical Bayes Methods. Technical Report 28, Stanford, Calif.: Division of Biostatistics, Stanford University, 1976. [p333]
- R. E. Tripp. *Non-Stochastic Ridge Regression and Effective Rank of the Regressors Matrix*. Ph.d. thesis, Department of Statistic, Virginia Polytechnic Institute and State University., 1983. [p326]
- W. N. Venables and B. D. Ripley. *Modern Applied Statistics with S*. Springer-Verlag, New York, 4th edition, 2002. URL <http://www.stats.ox.ac.uk/pub/MASS4>. ISBN 0-387-95457-0. [p326, 328, 333]
- H. D. Vinod. Application of New Ridge Regression Methods to a Study of Bell System Scale Economics. *Journal of the American Statistical Association*, 71(356):835–841, 1976. URL <http://doi.org/10.2307/2286847>. [p330, 332, 339]
- E. Wencheko. Estimation of the Signal-to-Noise in the Linear Regression Model. *Statistical Papers*, 41: 327–343, 2000. URL <https://doi.org/10.1007/BF02925926>. [p330]
- D. W. Wichern and G. A. Churchill. A Comparison of Ridge Estimators. *Technometrics*, 20(3):301–311, 1978. URL <http://doi.org/10.2307/1268139>. [p330]

Muhammad Imdad Ullah
Ph.D scholar (Statistics),
Department of Statistics
Bahauddin Zakariya University, Multan, Pakistan
ORCID ID: 0000-0002-1315-491X
mimdadasad@gmail.com

Muhammad Aslam
Associate Professor,
Department of Statistics
Bahauddin Zakariya University, Multan, Pakistan
aslamasadi@bzu.edu.pk

Saima Altaf
Assistant Professor,
Department of Statistics
Bahauddin Zakariya University, Multan, Pakistan
saimaaltaf@bzu.edu.pk